

Trichloro(1,1-dimethylbiguanidium- κN^3)zinc(II)

Miaoli Zhu,^a Liping Lu,^a Xianglin Jin^b and Pin Yang^{a*}

^aInstitute of Molecular Science, University of Shanxi, Taiyuan, Shanxi 030006, People's Republic of China, and ^bDepartment of Chemistry, University of Beijing, Beijing 100871, People's Republic of China

Correspondence e-mail: yangpin@sxu.edu.cn

Received 12 October 2001


Accepted 9 January 2002

Online 13 February 2002

The title compound, $[ZnCl_3(C_4H_{12}N_5)]$, was prepared from aqueous solution and its structure determined. The coordination geometry around the Zn atom is a tetrahedron, with the central Zn atom bound to three Cl atoms and to one N atom of the biguanide ligand. The dihedral angle between the two guanidine groups is $67.86(1)^\circ$.

Comment

An *N*-substituted derivative of biguanide, metformin (*N,N*-dimethylbiguanide), is a powerful oral antihyperglycaemic drug that has been used in many countries for over 30 years for treating diabetic patients with non-insulin-dependent diabetes mellitus. As an *N*-donor monodentate and bidentate ligand, it also forms complexes with many metal ions (Bailey, 1992; Ray, 1961). Zinc is considered an important inorganic chemical element involved in biological processes. It exists in many organs, such as the liver, kidney, brain and pancreas, and it has an important role and function within proteins, enzymes, and so on.



From a chemical point of view, biguanide and its *N*-substituted derivatives are considered to be a good type of chelating ligand, with abundant chemical properties. There are structural differences between monodentate and bidentate ligands when they coordinate to different metal ions, with the former normally forming a tetrahedron and the latter a square-planar configuration, e.g. the bidentate biguanide-derivative complexes of Co^{II} , Tc^V , Rh^V , Ag^{III} and Cu^{II} all have a square-planar geometry (Lemoine *et al.*, 1996; Marchi *et al.*, 1999;

Ghosh *et al.*, 1994; Hota & Saha, 1984). In addition, the planar angle between the two guanidine groups differs between complexes with monodentate or bidentate ligands. In the light of this, the title compound, (I), with a monodentate biguanide ligand, has been prepared, and its crystal structure is presented here.

The geometric parameters of (I) are listed in Table 1 and the molecular structure is illustrated in Fig. 1. The structure can be viewed as two parts, namely $C_4H_{12}N_5^+$ and $ZnCl_3^-$. These two parts are connected through the formation of a Zn–N coordination bond of 2.0611 (14) Å.

In the structure of (I), each Zn^{2+} cation is coordinated by one N atom of the biguanide ligand and three Cl^- ions, in a slightly distorted tetrahedral configuration. The Cl–Zn bond distances are almost equal; range 2.2568 (7)–2.2728 (6) Å, with a mean of 2.2622 Å. These values are compatible with the Co–Cl bond lengths in $[CoCl_3(C_4H_{12}N_5)]$ [2.250 (1)–2.264 (1) Å; Lemoine *et al.*, 1996]. The Cl–Zn distances in (I) are also in agreement with those found in a morpholine biguanide-coordinated Zn complex, $[ZnCl_3(C_6H_{14}N_5O)]$ (Yang & Zhu, 1991, 1992), in which the Cl–Zn bond distances [2.245 (1)–2.258 (1) Å] are slightly shorter than those in (I).

The Cl–Zn–Cl angles in (I) are in the range $109.82(3)$ – $116.50(3)^\circ$, and the Cl–Zn–N angles are in the range $104.43(4)$ – $108.45(4)^\circ$, close to the theoretical tetrahedral value of 109.5° .




Figure 1

A view of the molecular structure of (I) with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 2

A packing diagram of (I) viewed along the *b* axis.

In the crystal structure of metformin hydrochloride, Hariharan *et al.* (1989) pointed out that the C–N bond distances in the two guanide groups are intermediate between single and double bonds, in the range 1.330 (4)–1.348 (3) Å. A similar conclusion can be drawn for the C–N bond characterization in (I), with C–N bond lengths in the range 1.314 (2)–1.382 (2) Å. The standard single and double C–N bond lengths are 1.474 and 1.265 Å, respectively (Lide, 1962).

The two halves of the biguanide residue in (I) are planar, with atom C1 0.033 Å out of the N1/N2/N5 plane and atom C2 0.042 Å out of the N2/N3/N4 plane. The planes of the two different halves form a dihedral angle of 67.86° in (I); dihedral angles of 77.58 and 67.9 (1)° were found in [ZnCl₃(C₆H₁₄N₅O)] (Yang & Zhu, 1992) and metformin hydrochloride (Hariharan *et al.*, 1989), respectively. Thus, the dihedral angle in (I) is very close to that of the uncomplexed biguanide. It is noted that the different *N*-substituted group within the ligand is the main reason for the variation in the dihedral angle; the dihedral angle in the morpholine-substituted complex is approximately 9.72° larger than that in the dimethyl-substituted complex.

Experimental

Crystals of (I) were grown from an aqueous solution of zinc dichloride dihydrate (1.0 mmol) and *N,N*-dimethylbiguanide hydrochloride (1.0 mmol). The solution was left at room temperature and crystals formed in the bottom of the flask after 1–3 months. The elemental analyses were in agreement with the structural composition of (I).

Crystal data

[ZnCl₃(C₄H₁₂N₅)]
M_r = 301.91
 Monoclinic, *P*2₁/*c*
a = 12.5040 (7) Å
b = 7.5170 (5) Å
c = 13.0894 (5) Å
 β = 113.237 (3)°
V = 1130.5 (1) Å³
Z = 4

D_x = 1.774 Mg m⁻³
 Mo *K*α radiation
 Cell parameters from 10 407 reflections
 θ = 3.1–27.5°
 μ = 2.85 mm⁻¹
T = 173 (2) K
 Block, colourless
 0.4 × 0.3 × 0.2 mm

Data collection

Rigaku R-AXIS RAPID image-plate diffractometer
 Oscillation scans
 Absorption correction: empirical (ABSCOR; Higashi, 1995)
T_{min} = 0.402, *T_{max}* = 0.566
 10 407 measured reflections
 2574 independent reflections

2107 reflections with *I* > 2σ(*I*)
R_{int} = 0.029
 θ_{\max} = 27.5°
h = -16 → 16
k = -9 → 9
l = -16 → 15
 Intensity variation: ±1%

Refinement

Refinement on *F*²
R [*F*² > 2σ(*F*²)] = 0.022
wR (*F*²) = 0.048
S = 0.94
 2574 reflections
 138 parameters

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0243P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.56 \text{ e \AA}^{-3}$

H atoms attached to C atoms were treated as riding, with C–H = 0.96 Å and *U_{iso}*(H) = 1.5*U_{eq}* of the parent atom. H atoms on N atoms

Table 1

Selected geometric parameters (Å, °).

Zn1–N2	2.0611 (14)	N2–C2	1.382 (2)
Zn1–Cl3	2.2568 (7)	N3–C2	1.314 (2)
Zn1–Cl2	2.2570 (5)	N3–C4	1.459 (2)
Zn1–Cl4	2.2728 (6)	N3–C3	1.465 (2)
N1–C1	1.328 (2)	N4–C2	1.334 (2)
N2–C1	1.359 (2)	N5–C1	1.320 (2)
N2–Zn1–Cl3	107.05 (4)	C2–N3–C4	122.38 (15)
N2–Zn1–Cl2	108.45 (4)	C2–N3–C3	121.16 (16)
Cl3–Zn1–Cl2	116.50 (3)	C4–N3–C3	116.02 (15)
N2–Zn1–Cl4	104.43 (4)	N5–C1–N1	119.29 (16)
Cl3–Zn1–Cl4	109.82 (3)	N5–C1–N2	118.71 (16)
Cl2–Zn1–Cl4	109.86 (2)	N1–C1–N2	121.86 (15)
C1–N2–C2	118.68 (14)	N3–C2–N4	120.89 (16)
C1–N2–Zn1	121.78 (11)	N3–C2–N2	119.69 (15)
C2–N2–Zn1	117.15 (11)	N4–C2–N2	119.13 (15)
Zn1–N2–C1–N5	-8.2 (2)	C3–N3–C2–N4	-7.6 (3)
Zn1–N2–C1–N1	167.41 (13)	N3–C2–N2–C1	134.46 (17)
Zn1–N2–C2–N4	111.06 (16)	N1–C1–N2–C2	-30.7 (2)
Zn1–N2–C2–N3	-62.77 (19)	N5–C1–N2–C2	153.77 (16)
Cl4–Zn1–N2–C2	-107.20 (12)	N3–C2–N4–N2	173.8 (3)
Cl2–Zn1–N2–C2	9.91 (13)	N2–C1–N1–N5	-175.5 (3)
Cl3–Zn1–N2–C2	136.37 (11)		
C4–N3–C2–N4	164.46 (17)		

were refined with *U_{iso}*(H) = 0.08 Å² and N–H distances in the range 0.78–0.92 Å.

Data collection: *MSCIAFC Diffractometer Control Software* (Molecular Structure Corporation, 1994); cell refinement: *MSCIAFC Diffractometer Control Software*; data reduction: *SHELXS97* (Sheldrick, 1997); program(s) used to solve structure: *SHELXS97*; program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *XP* (Siemens, 1990).

We sincerely thank the National Natural Science Foundation of China and the Provincial National Foundation of Shanxi for support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: DE1178). Services for accessing these data are described at the back of the journal.

References

- Bailey, C. J. (1992). *Diabetes Care*, **15**, 755–772.
 Ghosh, S. P., Ghosh, M. C. & Gould, E. S. (1994). *Inorg. Chim. Acta*, **225**, 83–88.
 Hariharan, M., Rajan, S. S. & Srinivasan, R. (1989). *Acta Cryst.* **C45**, 911–913.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Hota, S. K. & Saha, C. R. (1984). *J. Coord. Chem.* **13**, 131–142.
 Lemoine, P., Chiadmi, M., Bissery, V., Tomas, A. & Viossat, B. (1996). *Acta Cryst.* **C52**, 1430–1436.
 Lide, D. R. Jr (1962). *Tetrahedron*, **17**, 125–134.
 Marchi, A., Marvelli, L., Cattabriga, M., Rossi, R., Neves, M., Bertolasi, V. & Ferrelli, V. (1999). *J. Chem. Soc. Dalton Trans.* pp. 1937–1943.
 Molecular Structure Corporation (1994). *MSCIAFC Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
 Ray, P. (1961). *Chem. Rev.* **61**, 313–359.
 Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
 Siemens (1990). *XP*. Version 4.2. Siemens Analytical X-ray Instruments Inc., Madison Wisconsin, USA.
 Yang, P. & Zhu, M. (1991). *Chin. Sci. Bull.* **36**, 1056.
 Yang, P. & Zhu, M.-L. (1992). *Chem. Res. Appl.* **4**, 20–25.