

*N,N*-Dimethylbiguanidium nitrate

Miaoli Zhu, Liping Lu and Pin Yang\*

Institute of Molecular Science, University of Shanxi, Taiyuan, Shanxi 030006, People's Republic of China

Correspondence e-mail: yangpin@sxu.edu.cn

## Key indicators

Single-crystal X-ray study

 $T = 184$  KMean  $\sigma(\text{C-N}) = 0.003$  Å $R$  factor = 0.054 $wR$  factor = 0.129


Data-to-parameter ratio = 10.1

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The cation in the title compound,  $\text{C}_4\text{H}_{12}\text{N}_5^+\cdot\text{NO}_3^-$ , is protonated at one imino group, and through this an intramolecular  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bond is formed, which stabilizes the conformation of the cation in the structure. The dihedral angle between the two guanidine groups is  $51.7(1)^\circ$ .

## Comment

Metformin is an antihyperglycemic agent which improves glucose tolerance in type-2 diabetic patients, lowering both basal and postprandial plasma glucose levels. There are some benefits for these diabetic patients who use it in order to control their plasma glucose levels. From pharmacological studies, metformin acts by improving peripheral sensitivity to insulin, inhibiting gastrointestinal absorption of glucose, and decreasing hepatic glucose production. Researchers indicate that metformin alone does not produce hypoglycemia in either diabetic or non-diabetic individuals (Davidson & Peters, 1997; Jackson *et al.*, 1987; Klip & Leiter, 1990). In previous reports, we studied that structures of complexes of metformin with  $\text{Zn}^{2+}$ ,  $\text{Cu}^{2+}$  and  $\text{Ni}^{2+}$  (Zhu *et al.*, 2002, 2002*a,b*). Magnesium, a ubiquitous element that plays a fundamental role in many cellular reactions, is involved in more than 300 enzymatic reactions in which food is catabolized and new chemical products are formed; these include glycogen breakdown, fat oxidation, protein synthesis, ATP synthesis, and the second messenger system (Lukaski, 2000). In order to obtain information regarding the interaction between metformin and magnesium ions, magnesium(II) nitrate was employed in our current research. However, the title compound, (I), was obtained instead of the  $\text{Mg}^{2+}$  complex. It can be compared with  $(\text{C}_4\text{H}_{12}\text{N}_5)[\text{TlBr}_4]$  (He *et al.*, 2002) and  $\text{C}_4\text{H}_{12}\text{N}_5^+\cdot\text{Cl}^-$  (Hariharan *et al.*, 1989).



Selected geometric parameters are listed in Table 1, and a perspective view of the structure is shown in Fig. 1. In the structure of (I), the planes of the two guanidine groups with a dihedral angle of  $51.7(1)^\circ$  planar angle. Atoms C1 and C2 deviate by  $-0.042(3)$  and  $-0.019(3)$  Å from the planes N1/N2/N3/C1 and N3/N4/N5/C2, respectively. The C–N bond distances of (I), in the range of  $1.324(3)$ – $1.336(3)$  Å, are shorter than a double bond, indicating a delocalization of

Received 11 March 2003

Accepted 24 March 2003

Online 31 March 2003




Figure 1

A view of the asymmetric unit of (I), with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radii. Dotted lines represent hydrogen bonds.




Figure 2

The dimer formed *via* hydrogen bonds, indicated by dashed lines.

$\pi$ -electron density across the biguanide group. 1,1-Dimethylbiguanide as a form of cation appears in several compounds, such as  $(C_4H_{12}N_5)[ZnCl_3]$  (Zhu *et al.*, 2002), and those previously mentioned. Compared with these compounds, a notable difference can be detected from the packing diagrams. As shown in Fig. 2, two molecules are connected by two  $N5-H\cdots N3$  hydrogen bonds, forming an elongated hexagon. The hydrogen bonding information is given in Table 2 and a packing diagram is shown in Fig. 3. In addition, the molecules in the crystal are held together by van der Waals forces and by a number of intermolecular  $N-H\cdots O$  interactions. A weak  $N-H\cdots N$  intramolecular hydrogen bond stabilizes the cation conformation (Fig. 1).

## Experimental

Crystals of (I) were grown from an aqueous solution of magnesium(II) nitrate hexahydrate (100.0 mmol) and *N,N*-dimethylbiguanide hydrochloride (100.0 mmol). The solution was left at room temperature and crystals formed after a few days. The elemental analysis result was in agreement with the structural composition of (I).

### Crystal data

$C_4H_{12}N_5^+ \cdot NO_3^-$	$Z = 2$
$M_r = 192.20$	$D_x = 1.418 \text{ Mg m}^{-3}$
Triclinic, $P\bar{1}$	Mo $K\alpha$ radiation
$a = 7.204$ (3) Å	Cell parameters from 2484 reflections
$b = 7.534$ (3) Å	$\theta = 2.8\text{--}24.7^\circ$
$c = 8.821$ (4) Å	$\mu = 0.12 \text{ mm}^{-1}$
$\alpha = 78.109$ (5)°	$T = 184$ (2) K
$\beta = 73.979$ (6)°	Block, colorless
$\gamma = 85.353$ (6)°	$0.30 \times 0.20 \times 0.10 \text{ mm}$
$V = 450.1$ (3) Å <sup>3</sup>	




Figure 3

A packing diagram of the structure of the title compound.

### Data collection

Bruker SMART 1K CCD area-detector diffractometer	1559 independent reflections
$\omega$ scans	1181 reflections with $I > 2\sigma(I)$
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)	$R_{\text{int}} = 0.022$
$T_{\text{min}} = 0.965$ , $T_{\text{max}} = 0.988$	$\theta_{\text{max}} = 25.0^\circ$
1875 measured reflections	$h = -5 \rightarrow 8$
	$k = -7 \rightarrow 8$
	$l = -10 \rightarrow 10$

### Refinement

Refinement on $F^2$	All H-atom parameters refined
$R[F^2 > 2\sigma(F^2)] = 0.054$	$w = 1/[\sigma^2(F_o^2) + (0.0627P)^2]$
$wR(F^2) = 0.129$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\text{max}} = 0.002$
1559 reflections	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$
155 parameters	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$

Table 1

Selected geometric parameters (Å, °).

C1—N1	1.327 (3)	C2—N5	1.324 (3)
C1—N2	1.335 (3)	C2—N3	1.331 (3)
C1—N3	1.343 (3)	C2—N4	1.336 (3)
N1—C1—N2	118.7 (2)	N3—C2—N4	124.1 (2)
N1—C1—N3	119.4 (2)	C1—N1—C4	121.3 (2)
N2—C1—N3	121.6 (2)	C1—N1—C3	121.2 (2)
N5—C2—N3	118.1 (2)	C2—N3—C1	121.5 (2)
N5—C2—N4	117.8 (2)	N2—C1—N1—C4	−164.5 (3)
N2—C1—N1—C4	−164.5 (3)	N5—C2—N3—C1	−159.1 (2)
N3—C1—N1—C4	21.6 (4)	N4—C2—N3—C1	23.8 (4)
N2—C1—N1—C3	6.3 (4)	N1—C1—N3—C2	−147.5 (2)
N3—C1—N1—C3	−167.6 (2)	N2—C1—N3—C2	38.8 (3)

Table 2

Hydrogen-bonding geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N5—H52 <sup>i</sup> ...O1 <sup>i</sup>	0.85 (3)	2.11 (3)	2.962 (3)	175 (3)
N5—H51 <sup>i</sup> ...N3 <sup>ii</sup>	0.91 (3)	2.10 (3)	3.003 (3)	171 (3)
N4—H41 <sup>i</sup> ...O3 <sup>i</sup>	0.84 (3)	2.12 (3)	2.932 (3)	162 (3)
N2—H21 <sup>i</sup> ...O1	0.87 (3)	2.13 (3)	2.980 (3)	166 (3)
N2—H22 <sup>i</sup> ...O1 <sup>iii</sup>	0.89 (3)	2.14 (3)	3.009 (3)	168 (3)
N4—H42 <sup>i</sup> ...O3 <sup>iv</sup>	0.90 (3)	2.10 (3)	2.936 (3)	153 (3)
N4—H42 <sup>i</sup> ...N2	0.90 (3)	2.58 (3)	2.913 (3)	103 (2)

Symmetry codes: (i)  $x, y, 1+z$ ; (ii)  $1-x, -y, 2-z$ ; (iii)  $1-x, 1-y, 1-z$ ; (iv)  $-x, 1-y, 1-z$ .

H atoms attached to C and N atoms were located in a difference Fourier map and refined with a global  $U_{\text{iso}}$  value. The C–H and N–H distances are in the ranges 0.93 (3)–0.98 (3) and 0.84 (3)–0.91 (3) Å, respectively.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SMART*; data reduction: *SAINTE* (Bruker, 2000); program(s) used to solve structure: *SHELXTL* (Sheldrick, 1999); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work is supported financially by the National Natural Science Foundation of China (No. 20171031 to PY), the Provincial Natural Foundation of Shanxi for Youth (No. 20011007 to LPL), and the Overseas Returned Scholar Foundation of Shanxi Province in 2002 for MLZ.

## References

- Bruker (2000). *SMART* (Version 5.0) and *SAINTE* (Version 6.02). Bruker AXS Inc., Madison, Wisconsin, USA.
- Davidson, M. B. & Peters, A. L. (1997). *Am. J. Med.* **102**, 99–110.
- Hariharan, M., Rajan, S. S. & Srinivasan, R. (1989). *Acta Cryst.* **C45**, 911–913.
- He, Z.-L., Zhu, M.-L. & Ma, G.-B. (2002). *Acta Cryst.* **E58**, m647–m649.
- Jackson, R. A., Hawa, M. I., Jaspan, J. B., Sim, B. M., Disilvio, L., Featherbe, D. & Kurtz, A. B. (1987). *Diabetes*, **36**, 632–640.
- Klip, A. & Leiter, L. A. (1990). *Diabetes Care*, **13**, 696–704.
- Lukaski, H. C. (2000). *Am. J. Clin. Nutr.* **72** (suppl.), 585S–593S.
- Sheldrick, G. M. (1999). *SHELXTL/PC*. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Zhu, M.-L., Lu, L.-P., Jin, X.-L. & Yang, P. (2002). *Acta Cryst.* **C58**, m158–m159.
- Zhu, M.-L., Lu, L.-P., Yang, P. & Jin, X.-L. (2002a). *Acta Cryst.* **E58**, m217–m219.
- Zhu, M.-L., Lu, L.-P., Yang, P. & Jin, X.-L. (2002b). *Acta Cryst.* **E58**, m272–m274.